































































- Tripathi, P., Dubey, N. K. & Shukla, A. K. (2008). Use of some essential oils as post- harvest botanical fungicides in the management of grey mould of grapes caused by *Botrytis cinerea*. *World Journal of Microbial Biotechnology*, 24, 39-46. <https://doi.org/10.1007/s11274-007-9435-2>.
- Umuhozariho, M. G., Shayo, N. B., Sallah, P. Y. K. & Msuya, J. M. (2013). Sensory evaluation of different preparation of cassava leaves from three species as a leafy vegetable. *African Journal of Biotechnology*, 13(46), 6452-6459. <https://doi.org/10.5897/AJB2013.12490>.
- Wijesundara, S. A. D. T. L., Kannangara, B. T. S. D. P. & Abeywickrama, K. (2015). Antifungal Activity of *Croton aromaticus* L. in vitro, Against Post- Harvest Fungal Pathogens Isolated from Tropical Fruits. *The Journal of Agricultural Sciences*, 11(2), 105-117. <https://doi.org/10.4038/jas.v11i2.8123>.
- Wilson, C. L., Solar, J. M., El Ghaouth, A. & Wisniewski, M. E. (1997). Rapid evaluation of plant extracts and essential oils for antifungal activity against *Botrytis cinerea*. *Plant Disease*, 81, 204-210. <https://doi.org/10.1094/PDIS.1997.81.2.204>.
- Zakaria, L., Sahak, S., Zakaria, M., & Salleh, B. (2009). Characterisation of *Colletotrichum* species associated with anthracnose of banana. *Tropical life sciences research*, 20(2), 119–125. PMID: 24575184; PMCID: PMC3819051. <https://pubmed.ncbi.nlm.nih.gov/24575184/>.
- Zhu, X., Wang, Q., Cao, J. & Jiang, W. (2008). Effects of chitosan coating on postharvest quality of mango (*Mangifera indica* L. cv. Tainong) fruits. *Journal of Food Process*. 32, 770–784. <https://doi.org/10.1111/j.1745-4549.2008.00213>.